

SUBMITTING A LETTER OF MEDICAL NECESSITY

A letter of medical necessity can provide a detailed rationale for why your patient needs a specific treatment, based on their medical history

You may provide a letter of medical necessity if:

- Your patient's claim was denied and you are submitting an appeal letter
- You are requesting a formulary exception or tiering exception to get access for your patient

SAFETY CONSIDERATIONS¹

Estrogen and progestin combinations, including ORIAHNN, increase the risk of thrombotic or thromboembolic disorders, including pulmonary embolism, deep vein thrombosis, stroke, and myocardial infarction, especially in women at increased risk for these events.

ORIAHNN is contraindicated in women with current or a history of thrombotic or thromboembolic disorders and in women at increased risk for these events, including women over 35 years of age who smoke and women with uncontrolled hypertension.

**HAVE A QUESTION?
CONTACT YOUR DEDICATED
ACCESS SPECIALIST**

**FOR SUPPORT OVER THE PHONE,
PLEASE CALL 1-800-ORIAHNN (1-800-674-2466)**

Please see additional Important Safety Information, including **BOXED WARNING** on THROMBOEMBOLIC AND VASCULAR EVENTS, on pages 4-5.

Please click here for full [Prescribing Information](#).

WRITING A LETTER OF MEDICAL NECESSITY

BEFORE YOU GET STARTED, MAKE SURE YOU HAVE THE FOLLOWING FOR AN EFFICIENT SUBMISSION OF YOUR LETTER OF MEDICAL NECESSITY:

- Patient's insurance policy/ID number
- Case ID number if a decision has already been rendered
- Patient's full name, plan identification number, and date of birth
- A brief medical history, including diagnosis, test results (ie, DXA bone density scan, if required), allergies, existing comorbidities, and International Classification of Diseases (ICD) code(s)
- Clinical support for your recommendation
- Your office contact information

THE LETTER OF MEDICAL NECESSITY MAY BE DOWNLOADED ONLINE OR COME FROM THE PAYER



Download and complete the online template from OriaHnnHCP.com/letter-of-medical-necessity-sample-template

OR



Complete the patient's payer version of the letter of medical necessity if available

As you navigate through the template, please fill in information based on your clinical assessment for your specific patient. Be sure to also include any other pertinent information for your patient.

Digital version available at
OriaHnnHCP.com/letter-of-medical-necessity-sample-template
Please open file on PC only to ensure usability

DXA, dual-energy X-ray absorptiometry.

SAFETY CONSIDERATIONS¹

ORIAHNN may cause a decrease in bone mineral density (BMD), which is greater with increasing duration of use and may not be completely reversible after stopping treatment. Assessment of BMD by dual-energy X-ray absorptiometry (DXA) is recommended at baseline and periodically thereafter. Limit the duration of use to 24 months to reduce the extent of bone loss.

This information is presented for informational purposes only and is not intended to provide reimbursement or legal advice. Providers are encouraged to contact third-party payers for specific information about their coverage policies. For more information, call an Access Specialist at 1-800-ORIAHNN (1-800-674-2466).

Please see additional Important Safety Information, including BOXED WARNING on THROMBOEMBOLIC AND VASCULAR EVENTS, on pages 4-5.

Please click [here](#) for full Prescribing Information.

 **OriaHnn™**
elagolix, estradiol and norethindrone acetate capsules
and elagolix capsules 300 mg/1 mg/0.5 mg and 300 mg

OriaHnn™ COMPLETE

SAMPLE LETTER OF MEDICAL NECESSITY



Before completing the template, ask the payer whether a specific form is required to help establish medical necessity.

[Date]
[Payer Name]
[Payer Address]
Attn: [Appeals Department]
Re: [Patient Name]
[Policy ID/Group Number]
[Date of Service]

To whom it may concern:

My name is [name] and I am a [board-certified medical specialty] [NPI] writing on behalf of my patient, [patient name], to request coverage for [product, dosage, and frequency]. [Patient Name] has been under my care for [X] months for the treatment of [disease or symptoms].

I am writing this letter for medical necessity because, after working with [patient name], I believe that [product name] is the best treatment for this patient, and it's important that a formulary exception be made.

[Provide a brief medical history, including diagnosis, allergies, existing comorbidities, and International Classification of Diseases (ICD) code(s)].

[Discuss rationale for using <product name> vs other treatments. Insert your recommendation summary here, including your professional opinion of your patient's likely prognosis or disease progression without treatment.]

[List of pertinent medical records] are enclosed, which offer additional support for the formulary exception request for [product name]. Please consider coverage of [product name] for my patient.

Please contact me at [telephone number] to answer any pending questions. I would be pleased to speak to the medical necessity of [product name] for [patient's name]'s [diagnosis].

Thank you in advance for your attention to this request.


Sincerely,

[Physician name and signature]
[Physician's medical specialty]
[Physician's NPI]
[Physician's practice name]
[Phone #]
[Fax #]

Please see Important Safety Information, including **BOXED WARNING** on **THROMBOEMBOLIC AND VASCULAR EVENTS**, on pages 4-5.

Please click [here](#) for full Prescribing Information.

Oriahnn[™] COMPLETE

 Oriahnn[™]
elagolix, estradiol and norethindrone acetate capsules
and elagolix capsules 300 mg/1 mg/0.5 mg and 300 mg

INDICATION AND IMPORTANT SAFETY INFORMATION FOR ORIAHNN™ (ELAGOLIX, ESTRADIOL, AND NORETHINDRONE ACETATE CAPSULES; ELAGOLIX CAPSULES)

INDICATION¹

ORIAHNN™ (elagolix, estradiol, and norethindrone acetate capsules; elagolix capsules) is indicated for the management of heavy menstrual bleeding associated with uterine leiomyomas (fibroids) in premenopausal women. Use of ORIAHNN should be limited to 24 months due to the risk of continued bone loss, which may not be reversible.

IMPORTANT SAFETY INFORMATION¹ THROMBOEMBOLIC AND VASCULAR EVENTS

Estrogen and progestin combinations, including ORIAHNN, increase the risk of thrombotic or thromboembolic disorders, including pulmonary embolism, deep vein thrombosis, stroke, and myocardial infarction, especially in women at increased risk for these events.

ORIAHNN is contraindicated in women with current or a history of thrombotic or thromboembolic disorders and in women at increased risk for these events, including women over 35 years of age who smoke and women with uncontrolled hypertension.

CONTRAINDICATIONS

- ORIAHNN is contraindicated in women at a high risk of arterial, venous thrombotic, or thromboembolic disorders; who are pregnant; with known osteoporosis; current or history of breast cancer or other hormonally sensitive malignancies; known hepatic impairment or disease; undiagnosed abnormal uterine bleeding; known anaphylactic reaction, angioedema, or hypersensitivity to ingredients of ORIAHNN; or with concomitant use of organic anion transporting polypeptide (OATP) 1B1 inhibitors that are known or expected to significantly increase elagolix plasma concentrations.

WARNINGS AND PRECAUTIONS

Thromboembolic Disorders and Vascular Events

- ORIAHNN is contraindicated in women with current or a history of thrombotic or thromboembolic disorders and in women at increased risk for these events. Components of ORIAHNN increase the risk of thrombotic or thromboembolic disorders, including pulmonary embolism, deep vein thrombosis, stroke, and myocardial infarction, especially in women at high risk for these events. In general, the risk is greatest among women over 35 years of age who smoke, and women with uncontrolled hypertension, dyslipidemia, vascular disease, or obesity.
- Discontinue ORIAHNN if an arterial or venous thrombotic, cardiovascular, or cerebrovascular event occurs. If feasible, discontinue ORIAHNN at least 4 to 6 weeks before surgery of the type associated with an increased risk of thromboembolism, or during periods of prolonged immobilization. Stop ORIAHNN

if there is sudden, unexplained partial or complete loss of vision, proptosis, diplopia, papilledema, or retinal vascular lesions and evaluate for retinal vein thrombosis immediately.

Bone Loss

- ORIAHNN is contraindicated in women with known osteoporosis. ORIAHNN may cause a decrease in bone mineral density (BMD) in some patients, which is greater with increasing duration of use and may not be completely reversible after stopping treatment.
- The impact of ORIAHNN-associated decreases in BMD on long-term bone health and future fracture risk is unknown. Consider the benefits and risks of ORIAHNN in patients with a history of low-trauma fracture or other risk factors for osteoporosis or bone loss, including those taking medications that may decrease BMD (e.g., systemic or chronic inhaled corticosteroids, anticonvulsants, or proton pump inhibitors).
- Assessment of BMD by dual-energy X-ray absorptiometry (DXA) is recommended at baseline and periodically thereafter. Consider discontinuing ORIAHNN if the risk associated with bone loss exceeds the potential benefit of treatment. Limit the duration of use to 24 months to reduce the extent of bone loss.

Hormonally Sensitive Malignancies

- ORIAHNN is contraindicated in women with current or a history of breast cancer and in women at increased risk for hormonally sensitive malignancies, such as those with mutations in BRCA genes.
- The use of estrogen alone and estrogen plus progestin has been reported to result in an increase in abnormal mammograms requiring further evaluation. Surveillance measures, such as breast examinations and regular mammography, are recommended. Discontinue ORIAHNN if a hormonally sensitive malignancy is diagnosed.

Suicidal Ideation, Suicidal Behavior, and Exacerbation of Mood Disorders

- Depression, depressed mood, and/or tearfulness were reported at a higher incidence in women taking ORIAHNN (3%) compared with placebo (1%) in the Phase 3 clinical trials. Suicidal ideation and behavior, including a completed suicide, occurred in women treated with lower doses of elagolix in clinical trials conducted for a different indication.
- Promptly evaluate patients with depressive symptoms to determine whether the risks of continued therapy outweigh the benefits. Patients with new or worsening depression, anxiety, or other mood changes should be referred to a mental health professional, as appropriate.

Please see additional Important Safety Information continued on page 5.

Please click here for full [Prescribing Information](#).



elagolix, estradiol and norethindrone acetate capsules
and elagolix capsules 300 mg/1 mg/0.5 mg and 300 mg

INDICATION AND IMPORTANT SAFETY INFORMATION FOR ORIAHNN™ (ELAGOLIX, ESTRADIOL, AND NORETHINDRONE ACETATE CAPSULES; ELAGOLIX CAPSULES) (CONT'D)

- Advise patients to seek immediate medical attention for suicidal ideation and behavior. Reevaluate the benefits and risks of continuing ORIAHNN if such events occur.

Hepatic Impairment and Transaminase Elevations

- ORIAHNN is contraindicated in women with known hepatic impairment or disease.
- Transaminase elevations in alanine aminotransferase (ALT) and aspartate aminotransferase (AST) occurred with ORIAHNN in Phase 3 clinical trials. No pattern in time to onset of these liver transaminase elevations was identified. Transaminase levels returned to baseline within 4 months after peak values in these patients.
- Instruct patients to promptly seek medical attention in case of symptoms or signs that may reflect liver injury, such as jaundice.

Elevated Blood Pressure

- ORIAHNN is contraindicated in women with uncontrolled hypertension. Maximum mean increases in systolic blood pressure occurred at Month 5, and a mean maximum increase in diastolic blood pressure occurred at Month 4 in ORIAHNN-treated women, as compared to placebo-treated women.
- For women with well-controlled hypertension, continue to monitor blood pressure and stop ORIAHNN if blood pressure rises significantly. Monitor blood pressure in normotensive women treated with ORIAHNN.

Gallbladder Disease or History of Cholestatic Jaundice

- Studies among estrogen users suggest a small increased relative risk of developing gallbladder disease. For women with a history of cholestatic jaundice associated with past estrogen use or with pregnancy, assess the risk-benefit of continuing therapy. Discontinue ORIAHNN if jaundice occurs.

Change in Menstrual Bleeding Pattern and Reduced Ability to Recognize Pregnancy

- ORIAHNN may delay the ability to recognize the occurrence of a pregnancy because it may reduce the intensity, duration, and amount of menstrual bleeding. Perform pregnancy testing if pregnancy is suspected and discontinue ORIAHNN if pregnancy is confirmed.
- The effect of hormonal contraceptives on the efficacy of ORIAHNN is unknown. Advise women to use non-hormonal contraception during treatment and for 1 week after discontinuing ORIAHNN.

Effects on Carbohydrate and Lipid Metabolism

- ORIAHNN may decrease glucose tolerance and result in increased glucose levels. More frequent monitoring in ORIAHNN-treated women with prediabetes and diabetes may be needed.
- In women with preexisting hypertriglyceridemia, estrogen therapy may be associated with elevations of plasma triglycerides leading to pancreatitis.

Use of elagolix is associated with increases in total cholesterol, low-density lipoprotein cholesterol (LDL-C), high-density lipoprotein cholesterol (HDL-C), and serum triglycerides. Monitor lipid levels and consider discontinuing ORIAHNN if hypercholesterolemia or hypertriglyceridemia worsens.

Alopecia

- In Phase 3 clinical trials, more women experienced alopecia, hair loss, and hair thinning with ORIAHNN (3.5%) compared to placebo (1.0%). In almost one-third of affected ORIAHNN-treated women, alopecia was the reason for discontinuing treatment. No specific pattern was described. In the majority of these women, hair loss was continuing when ORIAHNN was stopped. Whether the hair loss is reversible is unknown. Consider discontinuing ORIAHNN if hair loss becomes a concern.

Effect on Other Laboratory Results

- The use of estrogen and progestin combinations may raise serum concentrations of binding proteins (e.g., thyroid-binding globulin, corticosteroid-binding globulin), which may reduce the free thyroid or corticosteroid hormone levels. Patients with hypothyroidism and hypoadrenalism may require higher doses of thyroid hormone or cortisol replacement therapy, respectively.
- The use of estrogen and progestin may also affect the levels of sex hormone-binding globulin, coagulation factors, lipids, and glucose.

RISK OF ALLERGIC REACTIONS DUE TO THE INACTIVE INGREDIENT (FD&C YELLOW NO. 5)

- ORIAHNN contains FD&C Yellow No. 5 (tartrazine), which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. Although the overall incidence of FD&C Yellow No. 5 (tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin hypersensitivity.

ADVERSE REACTIONS

- Most common adverse reactions occurring in $\geq 5\%$ of women receiving ORIAHNN in clinical trials were hot flush, headache, fatigue, and metrorrhagia.

These are not all of the possible side effects of ORIAHNN.

Safety and effectiveness of ORIAHNN in pediatric patients have not been established.

Reference: 1. ORIAHNN [package insert]. North Chicago, IL: AbbVie Inc; 2020.

[Please click here for full Prescribing Information.](#)



Oriahnn™

elagolix, estradiol and norethindrone acetate capsules
and elagolix capsules 300 mg/1 mg/0.5 mg and 300 mg

abbvie

©2020 AbbVie Inc. North Chicago, IL 60064 US-GYNUF-200210 May 2020